



# CENTIPEDEGRASS LAWN CALENDAR

## Mowing Height:

1" - 2" raise 0.5" higher in hot weather. This is the most important practice for a healthy centipede grass lawn.

## Water:

1" per week if no rainfall.

## Fertilization: 1 - 2 lbs. N/1000 ft<sup>2</sup>/Yr

Follow fertilizer recommendations on soil test report. If the soil was not tested, use any turf fertilizer and follow label rates.

In the spring, do not apply nitrogen containing fertilizers until the soil temperature at the 4" depth is constantly 65°F and rising.

## Ideal pH Range: 5.0 - 6.0

Lime is rarely needed.

## Aeration:

Use a core aerator during active growth season.

**Dethatch:** If thatch exceeds 0.5" deep use a vertical mower with blades 1" apart; go over the lawn only one direction. Top dressing with 0.25" of soil can be effective.

## Seeding - New Lawn\*:

0.25 - 1.0 lb per 1,000ft<sup>2</sup>.

## Overseeding - Established Lawns\*:

Overseeding centipede grass with ryegrass is not recommended.

## Sodding:

500 ft<sup>2</sup> per pallet typical.

## Weed Control:

Read product label carefully to determine which weeds are controlled and on which grasses the product can be used.

**Spring preemergence\*** prevents crabgrass, goosegrass and other annual weeds.

**Broadleaf postemergence\*** spot spray to kill broadleaf plants like chickweed, wild violet, dandelion, wild onion, etc.

**Grassy weed postemergence\*** kills grassy weeds like crabgrass, dallisgrass, etc. Do not apply to drought-stressed centipede grass.

**Winter preemergence\*** prevents chickweed and other winter weed seeds from sprouting.

B		Best Month			P		Possible Month		M			Marginal Month	
Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec		
				OK	OK	OK	OK	OK					
			OK	OK	OK	OK	OK	OK					
				B	P	B	P						
				B	B	B	P						
				P	P	P							
				P	B	B	P						
			P	B	B	B	P	M	M				
	B	B	P										
OK	OK	OK	OK	OK	OK	OK	OK	OK	OK	OK	OK	OK	OK
				B	B	B	B	P					
								B	P				

\* Read weed control product labels carefully. Some products cannot be applied to lawns that will be seeded within a few weeks or when transitioning to active growth.